

Leading nucleon and the proton-nucleus inelasticity

J. Bellandi¹, J.R. Fleitas¹, and J. Dias de Deus²

¹*Instituto de Física, Universidade Estadual de Campinas, Campinas, SP 13083-970, Brazil*

²*Instituto Superior Técnico - CENTRA, Av. Rovisco Paes, 1, 1096 Lisboa Codex, Portugal*

Abstract

We present in this paper, a calculation of average proton-nucleus inelasticity. Using an interactive leading particle model and the Glauber model, we relate the leading particle distribution in nucleon-nucleus interactions with the respective one in nucleon-proton collisions. To describe the leading particle distribution in nucleon-proton, we use the Regge-Mueller formalism.

We calculate the average proton-nucleus inelasticity. Using an Iterative Leading Particle Model (Frichter et al., 1997) and the Glauber model (Glauber, 1959; Glauber et al., 1970), we relate the leading particle distribution in nucleon-nucleus interactions with the respective one in nucleon-proton collisions. In this model the leading particle spectrum in $p + A \rightarrow N(\text{nucleon}) + X$ collisions is built from successive interactions with ν interacting proton of the nucleus A and the behaviour is controlled by a straightforward convolution equation. It should be mentioned that, strictly speaking, the convolution should be 3-dimensional. Here we only considered the 1-dimension approximation. In a recent paper (Bellandi et al., 1999) we have used this model to describe the hadronic flux in the atmosphere, showing that the average nucleon-nucleus elasticity, $\langle x \rangle_{N-A}$, is correlated with the respective average nucleon-proton elasticity, $\langle x \rangle_{N-p}$, by means of the following relation

$$(1 - \langle x \rangle_{N-A}) = \frac{1}{\sigma_{in}^{N-A}} \int d^2b [1 - \exp[-(1 - \langle x \rangle_{N-p}) \sigma_{tot}^{pp} T(b)]] \quad (1)$$

where $T(b)$ is the nuclear thickness and given by means of the Woods-Saxon model (Woods, & Saxon, 1954; Barrett, & Jackson, 1977). Introducing the inelasticity given by $\langle k \rangle = 1 - \langle x \rangle$, this expression can be transformed in

$$\langle k \rangle_{N-A} = \frac{1}{\sigma_{in}^{N-A}} \int d^2b [1 - \exp[-\langle k \rangle_{N-p} \sigma_{tot}^{pp} T(b)]] . \quad (2)$$

It is clear from this relationship that only in small σ_{tot}^{pp} limit is $\langle K \rangle_{N-A} \simeq \langle K \rangle_N$. In general, $\langle K \rangle_{N-A} \geq \langle K \rangle_N$ and the effect increasing with the increase of σ_{tot}^{pp} . If $\langle K \rangle_N \rightarrow 0$, one also has $\langle K \rangle_{N-A} \rightarrow 0$. On the other hand, if $\langle K \rangle_N = 1$, then $\langle K \rangle_{N-A} = 1$, and Eq. (2) reproduces the Glauber model relationship between σ_{in}^{N-A} and σ_{tot}^{pp} . In order to calculate $\langle K \rangle_{N-A}$ we use for $\langle K \rangle_N$ the values calculated by means of the Regge-Mueller formalism (Batista et al., 1998) and as input for σ_{tot}^{pp} we have used the UA4/2 parametrizations for the energy dependence (Burnett et al., 1992). In the Fig. (1) we show the results of this calculations for the following nuclei: C, Al, Cu, Ag, Pb and air ($A = 14.5$). In this figure we also show recent experimental data for p-Pb, $\langle K \rangle = 0.84 \pm 0.16$ (Barroso et al., 1997) and for p-C, $\langle K \rangle = 0.65 \pm 0.08$ (Wilk & Włodarczyk, 1999).

In the Fig. (2), we compare the calculated $\langle K \rangle_{p-air}$ with results from some models used in Monte Carlo simulation (Gaisser et al., 1993); the Kopeliovich *et al.* (Kopeliovich et al., 1989) (KNP) QCD multiple Pomeron exchanges model; the Dual Parton model with sea-quark interaction of Capella *et al.* (Capella et al., 1981); the statistical model of Fowler *et al.* (Fowler et al., 1987) and with calculated values derived from cosmic ray data by Bellandi *et al.* (Bellandi et al., 1998). We note that the calculated $\langle K \rangle_{p-air}$ (Bellandi et al., 1998) was done assuming for the $T(b)$ nuclear thickness the Durand and Pi model (Durand & Pi, 1988), which gives small values for the average inelasticity. In the Fig. (2) we also show the average inelasticity values as calculated by means of this model.




Figure 1: Proton-nucleus inelasticities calculated. The up triangle is Pb data (Barroso et al., 1997) and down triangle is C data (Wilk & Włodarczyk, 1999) .




Figure 2: The $\langle K \rangle_{p-air}$ as a function of \sqrt{s} in GeV. The experimental data from (Bellandi et al., 1998). Dash line from (Fowler et al., 1985). Solid line from (Capella et al., 1981). Dot-dash line from (Kopeliovich et al., 1989). Dot line from Eq. (2) with Woods-Saxon model (Woods & Saxon, 1954; Barrett, & Jackson, 1977). Long-dash line from Durand-Pi model (Durand & Pi, 1988).

We would like to thank the Brazilian governmental agencies CNPq and CAPES for financial support

References

- Barrett, R.C. & Jackson, D.F. 1977, Nuclear Sizes and Structure, (Clarendon Press, Oxford).
- Barroso, S.L.C. *et al.* (Chacaltaya Collab.) 1997, Proc. 25th ICRC, (Durban, 1997)
- Batista, M. & Covolan, R.J.M. 1998, hep-ph:9811425.
- Bellandi, J., Fleitas, J.R. & Dias de Deus, J. 1998, Il Nuovo Cimento, **111A**, 149.
- Bellandi, J. *et al.* 1999, Proc. 26th ICRC, (Salt Lake City, 1999).
- Burnett, T.H. *et al.*, (UA4/2 Collab.) 1992, ApJ **349** L25.
- Capella, A. *et al.* 1981, Z. Phys. **C 10**, 249.
- Durand, L. & Pi, H. 1988, Phys. Rev. **D 38**, 78.
- Fowler, G. *et al.* 1987, Phys. Rev. **D 35**, 870.
- Frichter, G.M. *et al.* 1997, Phys. Rev. **D 56**, 3135.
- Gaisser, T.K. *et al.* 1993, Phys. Rev. **D 47**, 1919.
- Glauber, R.J. 1959, Lect. Theor. Phys. Vol.1, edited by W.Britten and L.G.Dunhan (Interscience, NY), 135.
- Glauber, R.J. *et al.* 1970, Nucl. Phys.**B 12**, 135.
- Kopeliovich, B.Z. *et al.* 1989, Phys. Rev. **D 39**, 769.
- Wilk, G. & Wlodarczyk, Z. 1999, Phys. Rev. **D 57**, 180.
- Woods, R.D. & Saxon, D.S. 1954, Phys. Rev. **95**, 577.
- Contribution to the 26th International Cosmic Ray Conference, Salt Lake City, Utah - August, 1999*